



ICTV Virus Taxonomy Profile: *Caliciviridae*

Jan Vinje^{1,*}, Mary K. Estes², Pedro Esteves³, Kim Y. Green⁴, Kazuhiko Katayama⁵, Nick J. Knowles⁶, Yvan L'Homme⁷, Vito Martella⁸, Harry Vennema⁹, Peter A. White¹⁰ and ICTV Report Consortium

Abstract

The family *Caliciviridae* includes viruses with single-stranded, positive-sense RNA genomes of 7.4–8.3 kb. The most clinically important representatives are human noroviruses, which are a leading cause of acute gastroenteritis in humans. Virions are non-enveloped with icosahedral symmetry. Members of seven genera infect mammals (*Lagovirus*, *Norovirus*, *Nebovirus*, *Recovirus*, *Sapovirus*, *Valovirus* and *Vesivirus*), members of two genera infect birds (*Bavovirus* and *Nacovirus*), and members of two genera infect fish (*Minovirus* and *Salovirus*). This is a summary of the International Committee on Taxonomy of Viruses (ICTV) Report on the family *Caliciviridae*, which is available at ictv.global/report/caliciviridae.

Table 1. Characteristics of members of the family *Caliciviridae*

Typical member:	Norwalk virus (M87661), species <i>Norwalk virus</i>, genus <i>Norovirus</i>
Virion	Non-enveloped with icosahedral symmetry, 27–40 nm in diameter
Genome	Single-stranded, positive-sense genomic RNA of 7.4–8.3 kb, with a 5'-terminal virus protein, genome-linked (VPg) and 3'-terminal poly(A)
Replication	Cytoplasmic
Translation	From genome-sized (non-structural proteins) and 3'-terminal subgenomic (structural proteins) mRNAs
Host range	Mammals (<i>Lagovirus</i> , <i>Norovirus</i> , <i>Nebovirus</i> , <i>Recovirus</i> , <i>Sapovirus</i> , <i>Valovirus</i> and <i>Vesivirus</i>), birds (<i>Bavovirus</i> , <i>Nacovirus</i>), fish (<i>Minovirus</i> , <i>Salovirus</i>)
Taxonomy	Realm <i>Riboviria</i> ; more than ten genera

VIRION

Calicivirus virions are 27–40 nm in diameter, non-enveloped with icosahedral symmetry (Table 1). The capsid is composed of 90 dimers of the major structural protein VP1 arranged on a T=3 icosahedral lattice (Fig. 1) [1]. Caliciviruses are characterised by a capsid architecture with 32 distinct cup-shaped depressions. Generally, caliciviruses are stable in the environment and enteric caliciviruses are acid-stable.

GENOME

Caliciviruses have a single-stranded, positive-sense genomic RNA of 6.4–8.5 kb organized into either two or three major ORFs, while a further ORF4 of murine norovirus encodes virulence factor 1 (VF1). A protein [virus protein, genome-linked (VPg), 10–15 kDa] is covalently linked to the 5'-terminus of genomic RNAs, which are also polyadenylated at their 3'-termini (Fig. 2). Genus-specific conserved nucleotide motifs are found

Received 04 September 2019; Accepted 05 September 2019; Published 01 October 2019

Author affiliations: ¹Division of Viral Diseases, National Center for Immunization and Respiratory Diseases, Centers for Disease Control and Prevention, Atlanta, GA, USA; ²Baylor College of Medicine, Houston, TX, USA; ³CIBIO/InBio – Centro de Investigação em Biodiversidade e Recursos Genéticos, Universidade do Porto, Campus de Vairão, 4485-661, Vairão, Portugal; ⁴Caliciviruses Section, Laboratory of Infectious Diseases, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, MD, USA; ⁵Laboratory of Viral Infection I, Kitasato Institute for Life Sciences Graduate School of Infection Control Sciences, Kitasato University, Tokyo, Japan; ⁶The Pirbright Institute, Pirbright, Surrey, UK; ⁷Université de Montréal, Montréal, Canada; ⁸Department of Veterinary Medicine, University Aldo Moro of Bari, Valenzano, Bari, Italy; ⁹Centre for Infectious Diseases Control, National Institute for Public Health and the Environment (RIVM), Bilthoven, The Netherlands; ¹⁰School of Biotechnology and Biomolecular Sciences, Faculty of Science, University of New South Wales, Sydney, Australia.

*Correspondence: Jan Vinje, jvinje@cdc.gov

Keywords: ICTV Report; taxonomy; *Caliciviridae*; norovirus.

Abbreviations: VPg, virus protein, genome-linked.

001332

This is an open-access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.




Fig. 1. The structure of the calicivirus capsid exemplified by a cryo-image reconstruction of recombinant Norwalk virus-like particles (left). X-ray structure of the Norwalk virus capsid (right) with the shell, protruding 1 and protruding 2 domains shown in blue, red and yellow, respectively. (Courtesy of B. V. Prasad.)




Fig. 2. Genome organization of human calicivirus MD-145 (AY032605, 7556 nt, genus *Norovirus*). Protein VPg is covalently linked to the 5'-end of genomic RNA and is depicted by a black circle. Cleavage sites in the ORF1 polyprotein are indicated by arrows; the flanking residues and amino acid coordinates are indicated, although these vary within the family. Pro: protease.

at the 5'-terminus of ORF1 and at the junction of the coding sequences for the non-structural/structural proteins.

REPLICATION

Replication of caliciviruses occurs in the cytoplasm in complexes on intracellular membranes by a VPg-mediated translation initiation process unique to the virus family and that uses genomic positive-sense RNA as the template to translate a large polyprotein that undergoes post-translational cleavage by a virus-encoded protease to form at least six mature non-structural proteins (NS1/2, NS3, NS4, NS5, NS6 and NS7) [2]. Subgenomic-sized, positive-sense RNA, co-terminal with the 3'-terminus of the genome, is the template for translation of VP1 as well as the 3'-terminal ORF product VP2 [3]. A dsRNA corresponding in size to full-length genomic RNA has been identified in feline calicivirus, murine norovirus and San Miguel sea lion virus-infected cells, indicating that replication occurs via a negative-sense intermediate. All caliciviruses require VPg; some require the function of eIF4E (feline calicivirus, porcine sapovirus), and some do not (murine norovirus).

PATHOGENICITY

Caliciviruses cause species-specific infections, with most noroviruses, sapoviruses and neboviruses restricted to the gastrointestinal tract; some lagoviruses, saloviruses and vesiviruses cause severe systemic infections in their natural hosts.

TAXONOMY

Viruses of seven genera (*Lagovirus*, *Norovirus*, *Nebovirus*, *Recovirus* [4], *Sapovirus*, *Valovirus* [5] and *Vesivirus*) infect a wide range of mammals, members of two genera infect birds (*Bavovirus* [6] and *Nacovirus* [7]) and members of two genera infect fish (*Minovirus* [8] and *Salovirus* [9]), while caliciviruses have also been detected in the greater green snake and frogs [10], highlighting the wide host range of viruses in the family. Caliciviruses are similar to picornaviruses in the presence of VPg and in sequence similarity of their RNA-directed RNA polymerase and protease proteins.

RESOURCES

Current ICTV Report on the family *Caliciviridae*: ictv.global/report/caliciviridae.

Funding information

Production of this summary, the online chapter, and associated resources was funded by a grant from the Wellcome Trust (WT108418AIA).

Acknowledgements

The findings and conclusions in this article are those of the authors and do not necessarily represent the official position of the Centers for Disease Control and Prevention. Members of the ICTV Report Consortium are Elliot J. Lefkowitz, Andrew J. Davison, Stuart G. Siddell, Peter Simmonds, Sead Sabanadzovic, Donald B. Smith, Richard J. Orton and Nick J. Knowles.

Conflicts of interest

The authors declare that there are no conflicts of interest.

References

- Prasad BVV, Hardy ME, Dokland T, Bella J, Rossmann MG. X-Ray crystallographic structure of the Norwalk virus capsid. *Science* 1999;286:287–290.
- Karst SM, Wobus CE, Goodfellow IG, Green KY, Virgin HW. Advances in norovirus biology. *Cell Host Microbe* 2014;15:668–680.
- Asanaka M, Atmar RL, Ruvolo V, Crawford SE, Neill FH *et al.* Replication and packaging of Norwalk virus RNA in cultured mammalian cells. *Proc Natl Acad Sci U S A* 2005;102:10327–10332.
- Farkas T, Sestak K, Wei C, Jiang X. Characterization of a rhesus monkey calicivirus representing a new genus of *Caliciviridae*. *J Virol* 2008;82:5408–5416.
- L'Homme Y, Sansregret R, Plante-Fortier Étienne, Lamontagne A-M, Ouardani M *et al.* Genomic characterization of swine caliciviruses representing a new genus of *Caliciviridae*. *Virus Genes* 2009;39:66–75.
- Wolf S, Reetz J, Otto P. Genetic characterization of a novel calicivirus from a chicken. *Arch Virol* 2011;156:1143–1150.
- Day JM, Ballard LL, Duke MV, Scheffler BE, Zsak L. Metagenomic analysis of the turkey gut RNA virus community. *Virol J* 2010;7:313.
- Mor SK, Phelps NBD, Ng TFF, Subramaniam K, Primus A *et al.* Genomic characterization of a novel calicivirus, FHMVCV-2012, from baitfish in the USA. *Arch Virol* 2017;162:3619–3627.
- Mikalsen AB, Nilsen P, Frøystad-Saugen M, Lindmo K, Eliassen TM *et al.* Characterization of a novel calicivirus causing systemic infection in atlantic salmon (*Salmo salar* L.): proposal for a new genus of *Caliciviridae*. *PLoS One* 2014;9:e107132.
- Shi M, Lin X-D, Chen X, Tian J-H, Chen L-J *et al.* The evolutionary history of vertebrate RNA viruses. *Nature* 2018;556:197–202.